

Morphological and molecular data to describe a hybrid population of the Common toad (*Bufo bufo*) and the Spined toad (*Bufo spinosus*) in western France

Tania Trujillo¹, Jorge Gutiérrez-Rodríguez¹, Jan W. Arntzen², Iñigo Martínez-Solano^{1,2,3,4,5,6}

¹ Museo Nacional de Ciencias Naturales, CSIC, c/ José Gutiérrez Abascal, 2, 28006 Madrid, Spain (present address)

² Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands

³ Instituto de Investigación en Recursos Cinegéticos (IREC-UCLM-CSIC-JCCM), Ronda de Toledo, s/n, 13005 Ciudad Real, Spain

⁴ CIBIO-InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Campus Agrário de Vairão, Universidade do Porto, 4485-661 Vairão, Portugal

⁵ Ecology, Evolution and Development Group, Department of Wetland Ecology, Doñana Biological Station, CSIC, c/ Americo Vespucio, s/n, 41092, Seville, Spain

⁶ Email: inigomsolano@mncn.csic.es

Key words: Amphibians, cross-amplification, genetic polymorphism, hybridization, microsatellites, morphology

Abstract

The use of hyper-variable markers across species is often hindered by low cross-species amplification success, a reduced level of polymorphism or a high frequency of null alleles. However, optimizing sets of reliable and informative markers that can be consistently amplified and scored across taxa is key to address questions about patterns of genetic diversity and structure, hybridization and speciation. Here we present 14 newly developed microsatellite markers in the Spined toad (*Bufo spinosus*), assess their polymorphism in two Iberian populations and test for cross-species amplification in the closely related Common toad (*Bufo bufo*). We then use the 12 loci co-amplifying in both species to the study of a morphologically intermediate population (Moyaux) from the contact zone in northwest France as well as reference populations of the two species from both sides of the contact zone. Individuals from Moyaux had mtDNA haplotypes of the two species and were identified as hybrids in analyses with software NewHybrids. These results provide solid evidence for ongoing hybridization between *B. bufo* and *B. spinosus*, with no apparent restrictions to gene flow.

Content

Introduction	1
Material and methods	2
Results	5
Discussion	7
Acknowledgements	8
References	8

Introduction

A widespread application of molecular markers across species is often hindered by a low cross-species ampli-

fication success, a high frequency of null alleles or a reduced level of polymorphism (Primmer *et al.*, 2005). This is particularly true for hyper-variable markers, like microsatellites, where high mutation rates are both advantageous (because their polymorphism allows very fine-scale temporal and spatial resolution) and problematic (because mutations in flanking regions reduce cross-amplification success) (Primmer and Merilä, 2002). Thus, optimizing sets of applicable, reliable and informative markers, which are key to address questions about patterns of genetic diversity and structure, hybridization and speciation, involves selecting highly polymorphic markers that can nonetheless be consistently amplified and scored in long diverged taxa.

Recent molecular studies have resolved the phylogeny of the *Bufo bufo* (Linnaeus, 1758) species group, revealing the existence of four species and delimiting their respective ranges (Litvinchuk *et al.*, 2008; García Porta *et al.*, 2012; Recuero *et al.*, 2012; Arntzen *et al.*, 2013a). As a result of these studies, *B. spinosus* Daudin, 1803 is defined as an Ibero-Maghrebian endemism, with populations in North Africa from Morocco to Tunisia, the Iberian Peninsula, the southern fringe of the British Isles (i.e., Jersey) and north of the Pyrenees across most of France, where it contacts *B. bufo* along a NW-SE line, roughly from Caen to Lyon (Arntzen *et al.*, 2013b, 2014). *Bufo bufo* is also present in the Apennine and Balkan peninsulas and extends as far east as northern Kazakhstan and eastern Siberia (Agasyan *et al.*, 2009).

Previous studies have used morphological characters, mtDNA and slowly evolving nuclear DNA markers to broadly delineate the contact zone between *B. bufo*



Fig. 1. Sampling localities of *Bufo bufo* (blue) and *B. spinosus* (red) in France: 1) Saumur (n=12); 2) Perré (n=8); 3) Durtal (n=8); 4) Moyaux (n=22); 5) Audresselles (n=14); 6) Erloy (n=20). The population at Moyaux (open circle) is identified as a *B. bufo* x *B. spinosus* hybrid population (see results). Black dots represent populations studied in Arntzen *et al.* (2013b), whereas the dashed line represents the contact zone as approximated in that study.

and *B. spinosus* in France (Arntzen *et al.*, 2013a, b, 2014). The results show an overall correspondence between morphology and molecules and congruence across molecular markers, with occasional discordance interpreted as resulting from incomplete lineage sorting, although hybridization-mediated gene flow cannot be ruled out.

Addressing questions about reproductive isolation versus hybridization and introgression in the two species requires characterization of molecular markers that amplify in both *B. bufo* and *B. spinosus* and the quantification of gene flow in areas of secondary contact. *Bufo bufo* has been the subject of a number of conservation genetics studies, especially in the UK and a number of polymorphic microsatellite loci are available (Scribner *et al.*, 1994, 1997, 2001; Hitchings and Beebee, 1998; Brede and Beebee, 2004, 2006; Wilkinson *et al.*, 2007). On the contrary, there are very few studies on *B. spinosus* and one of the reasons is low cross-amplification success of markers originally isolated from *B. bufo* (Brede *et al.*, 2001; Martínez-Solano and González, 2008; Arntzen *et al.*, 2014).

Here we describe a morphologically intermediate *Bufo* population from France with the help of a set of new polymorphic microsatellite markers isolated from a *B. spinosus* genomic library. We describe the variability of the newly developed markers in five populations of *B. spinosus* (two Iberian and three from west-

ern France), evaluate the cross-amplification success in two French populations of *B. bufo* and describe hybridization between the two species in a morphologically intermediate population in their contact zone.

Material and methods

A genomic library was generated from DNA of an adult female *B. spinosus* collected near Marbella, Málaga province, Spain, at the Sequencing Genotyping Facility, Cornell Life Sciences Core Laboratory Center, following the protocol described in Gutiérrez-Rodríguez *et al.* (2014) with minor modifications. Sequences containing microsatellites were scanned with Msatcommander (Faircloth, 2008) and sixty pairs of specific primers flanking regions that contained microsatellites were designed using the software Primerselect (DNASTAR, Inc.). Primers with an optimal melting temperature of 60°C were selected to facilitate multiplexing.

PCR reactions were performed using 5 × GoTaq Flexi buffer PROMEGA®; 3 mM MgCl₂; 0.3 mM dNTP; 0.3 μM of each primer; 0.5 U GoTaq Flexi DNA polymerase PROMEGA® and 1 μl of DNA, in a total volume of 15 μl. These initial tests included eight *B. spinosus* individuals from across the Iberian Peninsula and North Africa. PCR reactions consisted of initial denaturalization (95°C; 5 minutes), 40 cycles of

Table 1. Characterization of 14 new microsatellite loci in *Bufo spinosus* and polymorphism in two populations from Spain including locus designation, repeat motif, primer sequences, multiplex reaction, fluorescent dye, number of alleles (N_A) and observed (H_O) and expected (H_E) heterozygosity in the populations San Martín de la Vega/El Pardo (sample sizes: 94 and 32, respectively). An asterisk indicates a significant deviation from Hardy-Weinberg equilibrium associated with heterozygote deficit.

Locus	Repeat motif	Forward (top) and reverse (bottom) primers	Multiplex	Fluorescent dye	N_A	H_O	H_E
Bspi3.11	CTT	5' TTT CCT GCC TTC TTG TAA CGT TG 3' 5' CTC ACT GTC AGC AAT GTA TGA CC 3'	M1	6-FAM	6	0.4468/0.1250	0.4196/0.1190
Bspi4.24	GATA	5' AAA TTT GGG AGC AAT CCT GTA GG 3' 5' GTC ACA ACT GTC CTG TTA CCT TG 3'	M1	VIC	12	0.7234/0.7188	0.6821/0.7302
Bspi3.20	ACA	5' CCT CAG CAA ATA TAA GCC CAT GG 3' 5' TTG CAT CAG AAA CAA ACT TCC TG 3'	M1	NED	5	0.2043/0.3750*	0.6838/0.6066*
Bspi4.25	CTAT	5' ATT GTG TTG TGG ATG GAA CTA GC 3' 5' TAG AGA GAG CTG AAA TGT TGC TG 3'	M1	PET	12	0.6044/0.1667*	0.7873/0.8723*
Bspi4.16	GATA	5' AGG GCT ACA TAT CCT CTT CAG TG 3' 5' CTA AAC TGA GAA GAT GGC AAC CC 3'	M2	6-FAM	11	0.8085/0.8065	0.8277/0.7795
Bspi4.30	TCTA	5' CAC AGC CCT TTA CAA TCT ATC CG 3' 5' ACA AAC AGG CAG ACA AAT ATC AG 3'	M2	VIC	11	0.8191/0.7813	0.8474/0.8319
Bspi4.27	GATA	5' GAG ACA CGT AAT CCA GAC TTT CC 3' 5' TTA GGA CTT GTG TGA CAT CTG AG 3'	M2	NED	11	0.7979/0.6875	0.7742/0.6399
Bspi4.28	CTAT	5' GGC ATG GGT GAA TAA AGA AGT CC 3' 5' AAA CTT AGC TCA CCT GGT CAG C 3'	M2	PET	11	0.8830/0.8065	0.8593/0.7932
Bspi4.29	TAGA	5' CCC TTT CTA TGT CAC CTC TGT AC 3' 5' ACA GTG CCA TAT CTT CAG TGT TG 3'	M3	VIC	15	0.9149/0.9677	0.8498/0.8863
Bspi3.02	GGT	5' GGA TTA GGG CAT AGA CAA CTG AC 3' 5' CCG TCA CAG AGA AAT CAA AGG G 3'	M3	NED	3	0.2796/0.5333	0.3139/0.5859
Bspi3.19	ACT	5' CCG TCA CCA TTA CAA CTA CAC AG 3' 5' TTG TCA GAA GAA AGA GTG ATC GC 3'	M3	PET	5	0.3191/0.8438	0.5347/0.7450
Bspi3.26	AGT	5' GAA AGT CCC ATG TCT CGT TAT GC 3' 5' GTT GCC TCT TCT TCA CTC AGA TG 3'	M4	6-FAM	8	0.7447/0.8750	0.7148/0.7927
Bspi4.14	GATA	5' ATG AGT CTG CTA GGA ATT GTC TC 3' 5' CTG TAG CAA TCA TCT TCT CCT GC 3'	M4	VIC	12	0.8298/0.8710	0.8393/0.8096
Bspi4.07	GATA	5' ACC CAC CTC ACG TTA ATT ATT GC 3' 5' ATC TTC AAT TCC ATA GCT GCA CC 3'	M4	NED	9	0.8617/0.7097	0.8036/0.6716

denaturalization (95°C; 45 seconds), annealing (60°C; 45 seconds) and extension (72°C; 45 seconds) and final extension (72°C; 10 minutes). PCR products were visualized in 2% agarose gels to verify amplification and check the negative controls for potential contamination.

Fourteen loci that consistently amplified in all individuals and were polymorphic as seen on gel images were selected for further screening. These loci were amplified in four multiplex reactions designed with Multiplex Manager 1.0 (Holleley and Geerts, 2009). Forward primers were labeled with fluorescent dyes 6-FAM, VIC, NED and PET. Multiplex reactions were performed with Type-it Microsatellite PCR® (Qiagen) kits in a total volume of 15 µl that includes 7.5 µl of Master Mix, 1.2 µl of Primer Mix with 0.3 µM of each primer and 5.3 µl of RNase-free H₂O. Multiplex reactions consisted of initial denaturalization (95°C; 5

minutes), 30 cycles of denaturalization (95°C; 30 seconds), annealing (during 90 seconds at different temperatures in each multiplex reaction: multiplex reactions 1, 2 and 4 had an annealing temperature of 60°C and multiplex reaction 3 had a annealing temperature of 58°C) and extension (72°C; 30 seconds), and final extension (60°C; 30 minutes).

The above markers were tested in 126 *B. spinosus* tadpoles collected in two different populations in central Spain (Madrid province): 94 from San Martín de la Vega (40.23°, -3.59°) and 32 from El Pardo (40.56°, -3.95°). Additionally, we sampled individuals in six localities in northern France encompassing the northwestern end of the range to test cross-amplification in *B. bufo* and evaluate the utility of the markers to study hybridization with *B. spinosus*. Our sampling includes three populations in the “*spinosus*” side of the contact zone: Étang du Perré (n=8, 47.53°, -0.02°), Durtal (n=8,




Fig. 2. Top: individual assignment probabilities (P_A) in six French populations, based on NewHybrids results. Middle: individual scores of the first principal component of the genetic dataset (vertical axis), showing two groups consistent with species identification and with the population from Moyaux occupying an intermediate position. Bottom: P_A values for the 84 individuals genotyped based on Structure results for $K=2$. Red = *B. spinosus*; blue = *B. bufo*; grey = hybrid individuals. 1 = Saumur, 2 = Perré, 3 = Durtal, 4 = Moyaux, 5 = Audresselles and 6 = Erloy.

47.67°, -0.27°) and le Poitrineau, Saumur ($n=12$, 47.28°, -0.13°), two populations in the “*bufo*” side: Audresselles ($n=14$, 50.82°, 1.60°) and Autrepes (Erloy) ($n=20$, 49.92°, 3.83°) and a morphologically intermediate population (Moyaux, near Lisieux, $n=22$, 49.20°, 0.33°). Individuals from French populations were characterized with sequences of mtDNA and with diagnostic morphological characters as described in Arntzen *et al.* (2013b). DNA was extracted from approximately 0.25 g of tissue with Nucleospin® Tissue kits (Macherey-Nagel). Genotyping was performed on an ABI PRISM 3730 sequencer with the GeneScan 500 LIZ size standard (Applied Biosystems) and peaks were scored in GeneMapper 4.0 (Applied Biosystems).

Deviation from Hardy-Weinberg equilibrium and evidence of linkage disequilibrium across loci were tested for each population using Genepop 4.2 (Raymond and Rousset, 1995; Rousset, 2008). Using this software we also estimated the number of alleles (N_A),

observed (H_O) and expected heterozygosity (H_E) for each locus and population. Significance values for multiple tests were calculated applying a sequential Bonferroni correction (Rice, 1989). The presence of null alleles was tested using Micro-Checker 2.2.3 (Van Oosterhout *et al.*, 2004).

The software Genodive (Meirmans and Van Tien-deren, 2004) was used to perform a Principal Components Analysis (PCA) on the microsatellite genotype data in order to summarize multi-allelic information in a single vector. Individual scores of Principal Component 1 were then plotted against latitude to help visualize the power of discrimination of the newly developed markers in species identification.

We conducted analyses of the French dataset with the software Structure (Pritchard *et al.*, 2000). Runs consisted of a period of burn-in of 500,000 generations followed by 1,000,000 additional generations under an admixture model with allele frequencies corre-

Table 2. Data on polymorphism of twelve new microsatellite loci in French populations of *Bufo spinosus* (Perré, Durtal, Saumur) and *B. bufo* (Audresselles, Erloy) as well as the morphologically intermediate population Moyaux. Details include locus designation, observed size range, number of individuals (n), number of alleles (N_A), observed (H_O) and expected (H_E) heterozygosity.

Locality		Perré				Durtal				Saumur					
Locus	Size range	n	N_A	H_E	H_O	Size range	n	N_A	H_E	H_O	Size range	n	N_A	H_E	H_O
3.11	258-264	8	3	0.5417	0.5000	261-264	8	2	0.2333	0.2500	258-264	12	3	0.4746	0.5833
4.24	250-274	8	6	0.8500	1	246-278	8	6	0.8417	0.7500	250-270	12	4	0.7101	0.8333
4.25	188-204	8	4	0.6917	0.2500	160-204	6	5	0.8030	0.3333	188-208	12	5	0.8080	0.7500
4.16	226-250	8	5	0.8167	1	226-250	8	5	0.8417	0.7500	230-250	12	3	0.4674	0.5833
4.30	92-104	8	2	0.1250	0.1250	92-104	8	3	0.2417	0.2500	92-104	12	2	0.2283	0.2500
4.27	244-272	8	6	0.8333	0.8750	236-268	8	4	0.6750	0.7500	244-292	12	6	0.8442	1
4.28	185-205	8	5	0.7750	0.3750	185-213	8	4	0.3500	0.3750	185-201	12	3	0.5652	0.7500
4.29	202-258	8	10	0.9333	1	202-254	8	8	0.8917	0.8750	206-254	12	6	0.7717	0.5833
3.02	291-297	8	3	0.5417	0.7500	291-297	8	3	0.5917	0.5000	294-297	12	2	0.5181	0.5833
3.19	385-391	7	2	0.5275	0	373-394	5	4	0.8000	0	385	3	1	0	0
3.26	359-368	7	3	0.6264	0.5714	359	8	1	0	0	359-362	12	2	0.1594	0.1667
4.14	300-328	7	6	0.8462	0.8571	300-324	7	7	0.9000	0.8750	300-320	12	5	0.7971	1

Locality		Audresselles				Erloy				Moyaux					
Locus	Size range	n	N_A	H_E	H_O	Size range	n	N_A	H_E	H_O	Size range	n	N_A	H_E	H_O
3.11	264	14	1	0	0	264	20	1	0	0	258-264	21	3	0.3240	0.3333
4.24	226-266	13	6	0.7569	0.5385	226-258	12	5	0.6341	0.1667	226-274	20	9	0.8746	0.6000
4.25	188-212	14	7	0.8122	0.7857	180-212	20	8	0.8564	0.8000	184-216	21	8	0.8049	0.8095
4.16	218	14	1	0	0	218	20	1	0	0	218-250	22	6	0.5962	0.5454
4.30	92	14	1	0	0	92	20	1	0	0	92-104	22	3	0.4493	0.5909
4.27	254-298	14	12	0.9206	0.9286	252-302	20	13	0.9128	0.8500	246-290	21	13	0.9187	0.9524
4.28	187-233	14	9	0.9021	0.9286	183-233	20	9	0.8641	0.8500	185-233	22	9	0.8161	0.9091
4.29	212-256	14	10	0.8836	0.7143	206-252	20	13	0.9026	0.9000	202-246	22	15	0.9260	1
3.02	291	14	1	0	0	291	20	1	0	0	291-297	22	3	0.5613	0.6818
3.19	373-391	14	4	0.7540	0.8571	373-391	20	4	0.3872	0.4000	373-388	22	2	0.5106	0.4091
3.26	359-368	14	2	0.1376	0.1429	359-368	20	2	0.0974	0.1000	359-368	22	2	0.1691	0.1818
4.14	304-336	14	7	0.8518	0.9286	308-352	20	11	0.8628	0.9500	296-348	22	12	0.8837	0.9091

lated, for K values of 1 to 7 and 6 replicates per value of K. The optimal clustering scheme was selected through application of Evanno's criterion (Evanno *et al.*, 2005) as implemented in StructureHarvester (Earl and vonHoldt, 2012). Results were processed with the pipeline CLUMPAK (Kopelman *et al.*, 2015, available at: <http://clumpak.tau.ac.il/index.html>).

The French dataset was also analyzed with NewHybrids software (Anderson and Thompson, 2002) to estimate the posterior probability that genetically sampled individuals fall into each of a set of six categories (parental classes, F_1 , F_2 and the two backcrosses with the parental species). The latter four categories were subsequently combined into a hybrid score, which is calculated as the sum of the probabilities of assignment to each category.

In 22 adult toads from Moyaux (18 males and two

males and females from amplexed pairs) we studied a set of morphological characters that help to distinguish *B. bufo* from *B. spinosus*, namely Snout-Urostyle length (SUl), anterior and posterior parotoid distance (Pda, Pdp), length and width of the inner metatarsal tubercle (MTl, Mtw) and Paratoid angle (Pa) and the derived measurements Paratoid divergence (Pd=Pda/Pdp), Metatarsus Tubercle size (MTsize = MTl/SUl) and shape (MTshape = Mtw/MTl). For details on the measurements, data analysis and reference populations see Arntzen *et al.* (2013b).

Results

Fourteen loci consistently amplified and were polymorphic in the two populations of *B. spinosus* from




Fig. 3. Morphometric characteristics of adult toads from the population Moyaux (solid round symbols, 20 males, two females) relative to reference individuals from northern France, with ellipses showing one standard deviation around the mean, for *B. bufo* (shaded) and *B. spinosus* (unshaded). Reference data are taken from Arntzen *et al.* (2013b).

Spain (San Martín de la Vega and El Pardo). The original sequences were deposited in GenBank under accession numbers KX271337-KX271350.

The number of observed alleles per locus across the 14 loci optimized in *B. spinosus* ranged from 2 to 14 (mean $N_A=8.1$) in San Martín de la Vega and from 2 to 12 (mean $N_A=6.4$) in El Pardo. Observed and expected heterozygosities ranged from 0.20 to 0.91 (mean $H_O=0.66$) and 0.31 to 0.86 (mean $H_E=0.71$) respectively in San Martín de la Vega; and from 0.13 to 0.97 (mean $H_O=0.66$) and 0.12 to 0.89 (mean $H_E=0.70$) in El Pardo. Micro-Checker inferred the likely presence of null alleles in loci Bspi3.19, Bspi3.20 and Bspi4.25, associated with deviations from Hardy Weinberg equilibrium due to heterozygote deficits in both populations (Bspi3.20 and 4.25) and San Martín de la Vega

(Bspi3.19); therefore, caution should be taken when using these markers for population studies in *B. spinosus*. Four locus x locus combinations significantly deviated from linkage disequilibrium (after accounting for the Bonferroni correction, $P<0.001$) in Iberian populations (two in each population, with no consistent pattern across populations).

Locus Bspi4.07 only amplified in *B. spinosus* and locus Bspi3.20 did not consistently amplify in *B. bufo*. Therefore, we calculated the number of alleles (N_A), observed (H_O) and expected (H_E) heterozygosity in the remaining populations using twelve microsatellites: Bspi3.11, Bspi4.24, Bspi4.25, Bspi4.16, Bspi4.30, Bspi4.27, Bspi4.28, Bspi4.29, Bspi3.02, Bspi3.19, Bspi3.26 and Bspi4.14. Results are shown in Table 2.

Loci Bspi4.16, 3.02, 3.11 and 4.30 were monomor-

phic in the two *B. bufo* populations analyzed. Significant deviations from Hardy-Weinberg equilibrium (H_i : heterozygote deficit) were detected for loci Bspi3.19 in Durtal, Bspi4.28 in Perré and Bspi4.24 in Moyaux and Erloy. Significant linkage disequilibrium were only observed in Moyaux (four locus \times locus combinations: Bspi4.24 \times Bspi4.27, Bspi4.24 \times Bspi4.29, Bspi4.28 \times Bspi4.29 and Bspi4.28 \times Bspi4.14).

All individuals analyzed from Perré (n=8), Durtal (n=5) and Saumur (n=10) had the characteristic mtDNA profile of *B. spinosus*, whereas all individuals from Audresselles (n=22) and Erloy (n=28) had *B. bufo* mtDNA. In Moyaux, six individuals had *B. spinosus* mtDNA and 16 individuals had *B. bufo* mtDNA haplotypes. We found two interspecific amplexed pairs in our sample, a male *B. spinosus* mating with a female *B. bufo* and a male *B. bufo* with a female *B. spinosus*.

The first principal component of the microsatellite genotype data explained 25.8% of variation; when individual scores were plotted against geography, a pattern emerged of three groups corresponding to 1) the populations of Saumur, Durtal and Perré; 2) Moyaux and 3) Audresselles and Erloy (Fig. 2).

Structure analyses grouped individuals in two clusters (Fig. 2), with high assignment probabilities. A first cluster, corresponding to *B. spinosus*, included the populations Saumur (average probability of assignment, P_A : 0.99), Perré (P_A : 0.97), Durtal (P_A : 0.99), whereas the second cluster (*B. bufo*) included the populations Erloy (P_A : 0.99) and Audresselles (P_A : 0.99). Individuals from Moyaux were generally admixed, with a higher probability of assignment to the *B. bufo* cluster (P_A : 0.85, range 0.54-0.98, Fig. 2).

According to the results of NewHybrids analyses (Fig. 2) most individuals in the populations of Saumur, Perré and Durtal correspond to *B. spinosus* (population average proportions of 1.00, 0.63 and 0.88 respectively), while the remainder of the individuals from these populations (three individuals in Perré and one individual in Durtal) was assigned to the F_2 class. In the populations Audresselles and Erloy all individuals were classified as *B. bufo* with high probability, whereas in Moyaux, most individuals (64%) were classified as backcrosses with *B. bufo*, 27% were classified as F_2 individuals and the remaining 9% individuals were classified as *B. bufo* (Fig. 2).

The morphometric characteristics of the population Moyaux were intermediate between those of *B. bufo* and *B. spinosus*, as shown by the analysis of the diagnostic characters paratoid angle / paratoid divergence and metatarsus tubercle shape and size (Fig. 3).

Discussion

The new set of 14 polymorphic microsatellite markers is a valuable tool for population studies on *B. spinosus*. The species, formerly abundant throughout its Iberian range, is now undergoing a slow but sustained decline (Ortiz Santaliestra, 2014), and molecular tools can provide relevant information about patterns of genetic diversity and structure to guide conservation efforts. Furthermore, the high cross-amplification rates and observed levels of polymorphism in *B. bufo* make these markers valuable for fine-scale studies of the contact zone with *B. spinosus* as well as to complement previously developed markers in local or regional studies on *B. bufo*. Additional populations of the latter species should be assayed with the new markers to better characterize actual polymorphism, which was certainly underestimated in this study (for instance, markers Bspi3.02, 3.11 and 4.30 are monomorphic in the two French populations of *B. bufo* analyzed here, but additional polymorphism was found in a reduced sample of eastern European samples, data not shown).

Based on the new molecular data, the morphologically intermediate population of Moyaux is shown to represent a hybrid population, where mtDNA haplotypes of the two species can be found in syntopy. Co-occurrence of mtDNA haplotypes of the two species in a single site is rare, but had been previously reported (Recuero *et al.*, 2012; Arntzen *et al.*, 2013a). However, microsatellites also reveal a clearly admixed nuclear DNA genetic pool at Moyaux, suggestive of interspecific gene flow. This is thus the first solid evidence of hybridization between the two species (see Arntzen *et al.*, 2013a for a discussion about alternative interpretations of allozyme data on the taxonomic status of *B. bufo* and *B. spinosus*). While more detailed transects need to be studied to better characterize the contact zone in terms of its relative width or patterns of symmetry/asymmetry of gene flow across species, our preliminary results show a consistent signal of hybridization across independent sets of markers, including species-diagnostic morphological characters, mtDNA haplotypes and microsatellite loci. Thus, at least some of the previously observed instances of nuclear DNA haplotype sharing in slowly evolving nuclear markers (POMC, RAG1) may also result from interspecific gene flow rather than from incomplete lineage sorting alone (Arntzen *et al.*, 2013a, 2014).

The lack of F_1 hybrids and the high frequency of F_2 and backcrossed individuals in the hybrid population

at Moyaux suggest no restrictions to gene flow (random mating), which is in accordance with the observation of amplexant mates with mtDNA from the two different species (in both directions). This contrasts with the high incidence of deviations from linkage disequilibrium in the hybrid population, which may be indicative of assortative mating or selection rather than result from random demographic effects, because hybridization tends to break down species-specific physical associations between markers in hybrid zones and create new ones (Jiggins and Mallet, 2000). In any case, the high frequency of admixed individuals and relative scarcity of representatives of the parental taxa indicates this would be a relatively old contact zone, with many generations of hybrids gradually increasing their frequency in the population and becoming the dominant parental class, although more detailed, replicated transects are required to learn more about the potential role of selection in shaping the contact zone.

Whereas the distinction of *B. spinosus* as a separate species from *B. bufo* has long been subject to confusion, close inspection of the accumulating morphological and molecular evidence for species differentiation indicates that the two species have diagnostic morphological features and are highly distinct in their mitochondrial and nuclear DNA, except, as shown here, in parts of the contact zone where some hybridization occurs (including some of the reference *B. spinosus* populations, which according to NewHybrids results may also include some hybrid individuals, see Fig. 2). More detailed analyses of different sections of this contact zone are required to further delineate it and investigate whether observed patterns are generalizable. Additionally, comparative data about demographic and life history traits across species may be relevant to understand the evolutionary processes maintaining species boundaries in the face of gene flow. For instance, the skeletochronological work of Hemelaar (1988) showed remarkable differences in age at maturity and growth rates in *Bufo* populations from different latitudes. Of the five populations compared, one from southern France represents *B. spinosus* as inferred from recent molecular studies, whereas the other four would correspond to *B. bufo* on geographic grounds. Several characteristics of the French population stand out in comparison with populations of *B. bufo*, for instance much faster growth rates, larger maximum body size and earlier age at maturity, suggesting genetically, rather than environmentally induced differences (Hemelaar, 1988). Also, *B. spinosus* appears to start breeding a few weeks earlier than *B. bufo* and the

breeding season is also more prolonged than in *B. bufo*, which is usually considered an explosive breeder. These differences, in concert with potential mechanisms of species recognition, which do not seem to play a role in preventing interspecific mating in Moyaux but may be apparent elsewhere, can shape patterns of interspecific gene flow in the contact zone, ultimately determining its evolutionary fate.

Acknowledgements

The authors thank G. Rodríguez, G. Sánchez-Montes, A. Vallvé, J. van Alphen, A. Zuiderwijk and Agentes Forestales de la Comunidad de Madrid for help in collecting individuals and P. Ochoa for technical support. Two anonymous reviewers provided valuable feedback on a previous draft. This research was funded by grants CGL2008-04271-C02-01/BOS and CGL2011-28300 (Ministerio de Ciencia e Innovación), Ministerio de Economía y Competitividad and FEDER) and PII10-0097-4200 (Junta de Comunidades de Castilla la Mancha) to IMS. J. Gutiérrez-Rodríguez was supported by the Consejo Superior de Investigaciones Científicas of Spain (CSIC) and the European Social Fund (ESF) with a JAE-pre PhD fellowship. IMS was funded by the project 'Biodiversity, Ecology and Global Change', co-financed by North Portugal Regional Operational Programme 2007/2013 (ON.2–O Novo Norte), under the National Strategic Reference Framework (NSRF), through the European Regional Development Fund (ERDF) and by funding from the Spanish Severo Ochoa Program (SEV-2012-0262). IMS's stay in Leiden was supported by a Naturalis Temminck fellowship.

References

- Agasyan A, Avisi A, Tuniyev B, Crnobrnja Isailovic J, Lymberakis P, Andrén C, Cogalniceanu D, Wilkinson J, Ananjeva N, Üzümlü N, Orlov N, Podlucky R, Tuniyev S, Kaya U. 2009. *Bufo bufo*. The IUCN Red List of Threatened Species. Version 2014.2. [<http://www.iucnredlist.org>]
- Anderson EC, Thompson EA. 2002. A model-based method for identifying species hybrids using multilocus genetic data. *Genetics* 160: 1217-1229.
- Arntzen JW, Recuero E, Canestrelli D, Martínez-Solano I. 2013a. How complex is the *Bufo bufo* species group? *Molecular Phylogenetics and Evolution* 69: 1203-1208.
- Arntzen JW, McAtear J, Recuero E, Ziermann JM, Ohler A, van Alphen J, Martínez-Solano I. 2013b. Morphological and genetic differentiation of *Bufo* toads: two cryptic species in Western Europe (Anura, Bufonidae). *Contributions to Zoology* 82: 147-169.
- Arntzen JW, Wilkinson JW, Butôt R, Martínez-Solano I. 2014. A new vertebrate species native to the British Isles: *Bufo spinosus* Daudin, 1803 in Jersey. *Herpetological Journal* 24: 209-216.
- Brede E, Rowe G, Trojanowski J, Beebe TJC. 2001. Polymerase chain reaction primers for microsatellite loci in the

- Common Toad *Bufo bufo*. *Molecular Ecology Notes* 1: 308-310.
- Brede EG, Beebee TJC. 2004. Contrasting population structures in two sympatric anurans: implications for species conservation. *Heredity* 92: 110-117.
- Brede EG, Beebee TJC. 2006. Large variations in the ratio of effective breeding and census population sizes between two species of pond-breeding anurans. *Biological Journal of the Linnean Society* 89: 365-372.
- Daudin FM. 1803. Histoire Naturelle, Générale et Particulière des Reptiles; Ouvrage Faisant suite à l'Histoire Naturelle Générale et Particulière, Composée par Leclerc de Buffon; et Rédigée par C.S. Sonnini, Membre de Plusieurs Sociétés Savantes. Volume 8. Paris: F. Dufart.
- Earl DA, vonHoldt BM. 2012. *Structure Harvester*: a website and program for visualizing *Structure* output and implementing the Evanno method. *Conservation Genetics Resources* 4: 359-361.
- Evanno G, Regnaut S, Goudet J. 2005. Detecting the number of clusters of individuals using the software *Structure*: a simulation study. *Molecular Ecology* 14: 2611-2620.
- Faircloth BC. 2008. *Msatcommander*: detection of microsatellite repeat arrays and automated, locus-specific primer design. *Molecular Ecology Resources* 8: 92-94.
- García-Porta J, Litvinchuk SN, Crochet PA, Romano A, Geniez PH, Lo-Valvo M, Lymberakis P, Carranza S. 2012. Molecular phylogenetics and historical biogeography of the west-palaearctic common toads (*Bufo* species complex). *Molecular Phylogenetics and Evolution* 63: 113-130.
- Gutiérrez Rodríguez J, Salvi D, Geffen E, Gafny S, Martínez-Solano I. 2014. Isolation and characterisation of novel polymorphic microsatellite loci in Iberian painted frogs (*Discoglossus galganoi* and *D. jeanneae*), with data on cross-species amplification in *Discoglossus* and *Latonia* (Alytidae). *Herpetological Journal* 24: 261-265.
- Hemelaar A. 1988. Age, growth and other population characteristics of *Bufo bufo* from different latitudes and altitudes. *Journal of Herpetology* 22: 369-388.
- Hitchings SP, Beebee TJC. 1998. Loss of genetic diversity and fitness in common toad (*Bufo bufo*) populations isolated by inimical habitat. *Journal of Evolutionary Biology* 11: 269-283.
- Holley CE, Geerts PG. 2009. *Multiplex Manager* 1.0: a cross-platform computer program that plans and optimizes multiplex PCR. *BioTechniques* 46: 511-517.
- Jiggins CD, Mallet J. 2000. Bimodal hybrid zones and speciation. *Trends in Ecology and Evolution* 15: 250-255.
- Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I. 2015. *Clumpak*: a program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources* 15: 1179-1191.
- Linnaeus C. 1758. *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis*. 10th Edition. Volume 1. Stockholm: L. Salvii.
- Litvinchuk S, Borkin L, Skorinov D, Rosanov J. 2008. A new species of common toads from the Talysh mountains, south-eastern Caucasus: genome size, allozyme, and morphological evidences. *Russian Journal of Herpetology* 15: 19-43.
- Martínez-Solano I, González EG. 2008. Patterns of gene flow and source-sink dynamics in high altitude populations of the common toad *Bufo bufo* (Anura: Bufonidae). *Biological Journal of the Linnean Society* 95: 824-839.
- Meirmans PG, Van Tienderen PH. 2004. *Genotype and Genodive*: two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes* 4: 792-794.
- Ortiz Santaliestra ME. 2014. Sapo común – *Bufo spinosus*. In: *Enciclopedia Virtual de los Vertebrados Españoles*. A. Salvador & I. Martínez-Solano (Eds.). Madrid: Museo Nacional de Ciencias Naturales. [<http://www.vertebradosibericos.org/>]
- Primmer CR, Merilä J. 2002. A low rate of cross-species microsatellite amplification success in Ranid frogs. *Conservation Genetics* 3: 445-449.
- Primmer CR, Painter JN, Koskinen MT, Palo JU, Merilä J. 2005. Factors affecting avian cross-species microsatellite amplification. *Journal of Avian Biology* 36: 348-360.
- Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155: 945-959.
- Raymond M, Rousset F. 1995. *Genepop* (version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity* 86: 248-249.
- Recuero E, Canestrelli D, Vörös J, Szabó K, Poyarkov NA, Arntzen JW, Crnobrnja-Isailovic J, Kidov AA, Cogălniceanu D, Caputo FP, Nascetti G, Martínez-Solano I. 2012. Multilocus species tree analyses resolve the radiation of the widespread *Bufo bufo* species group (Anura, Bufonidae). *Molecular Phylogenetics and Evolution* 62: 71-86.
- Rice WR. 1989. Analyzing tables of statistical tests. *Evolution* 43: 223-225.
- Rousset F. 2008. *Genepop'007*: a complete reimplementation of the *Genepop* software for Windows and Linux. *Molecular Ecology Resources* 8: 103-106.
- Scribner KT, Arntzen JW, Burke T. 1994. Comparative analysis of intra- and interpopulation genetic diversity in *Bufo bufo*, using allozyme, single-locus microsatellite, minisatellite, and multilocus minisatellite data. *Molecular Biology and Evolution* 11: 737-748.
- Scribner KT, Arntzen JW, Burke T. 1997. Effective number of breeding adults in *Bufo bufo* estimated from age-specific variation at minisatellite loci. *Molecular Ecology* 6: 701-712.
- Scribner KT, Arntzen JW, Cruddace N, Oldham R, Burke T. 2001. Environmental correlates of toad abundance and population genetic diversity. *Biological Conservation* 98: 201-210.
- van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. 2004. *Micro-Checker*: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4: 535-538.
- Wilkinson JW, Beebee TJC, Griffiths RA. 2007. Conservation genetics of an island toad: *Bufo bufo* in Jersey. *Herpetological Journal* 17: 192-198.

Received: 12 January 2016

Revised and accepted: 15 June 2016

Published online: 14 February 2017

Editor: A. Ivanović

